arXiv:1305.0025v1 [gr-qc] 30 Apr 2013

Einstein static Universe in hybrid metric-Palatini gravity

Christian G. Böhmer[1],* Francisco S.N. Lobo[2],† and Nicola Tamanini[1]‡
[1]*Department of Mathematics and Institute of Origins,*
University College London, Gower Street, London, WC1E 6BT, UK and
[2]*Centro de Astronomia e Astrofísica da Universidade de Lisboa,*
Campo Grande, Ed. C8 1749-016 Lisboa, Portugal
(Dated: May 2, 2013)

Hybrid metric-Palatini gravity is a recent and novel approach to modified theories of gravity, which consists of adding to the metric Einstein-Hilbert Lagrangian an $f(\mathcal{R})$ term constructed à la Palatini. It was shown that the theory passes local tests even if the scalar field is very light, and thus implies the existence of a long-range scalar field, which is able to modify the dynamics in galactic and cosmological scales, but leaves the Solar System unaffected. In this work, motivated by the possibility that the Universe may have started out in an asymptotically Einstein static state in the inflationary universe context, we analyse the stability of the Einstein static Universe by considering linear homogeneous perturbations in the respective dynamically equivalent scalar-tensor representation of hybrid metric-Palatini gravity. Considering linear homogeneous perturbations, the stability regions of the Einstein static universe are parametrized by the first and second derivatives of the scalar potential, and it is explicitly shown that a large class of stable solutions exists in the respective parameter space, in the context of hybrid metric-Palatini gravity.

PACS numbers: 04.20.Jb, 04.50.Kd

I. INTRODUCTION

The standard model of cosmology is remarkably successful in accounting for the observed features of the Universe and during the last two decades has evolved from being mainly a theoretical area of Physics to become a field supported by observational data of high-precision. Recent experiments call upon state of the art technology in Astronomy and in Astrophysics to provide detailed information about the contents and history of the Universe. This has led to the measuring of the parameters that describe our Universe with increasing precision. In this context, the Nobel Prize of Physics 2011 was attributed for the discovery of the accelerating expansion of the Universe through observations of distant supernovae [1]. In fact, after fifteen years of extensive research, we still lack a fundamental understanding of this late-time cosmic acceleration, and its resolution will be one of the major objectives in cosmology during the next decade with several surveys and experiments to address the nature of the cosmic speed-up. Recently, the released Planck data of the 2.7 full sky survey [2] have also shown a number of intriguing features, which will require a change in our standard view of the Universe.

The standard model of cosmology has favoured a missing stress-energy component, in particular, the dark energy models [3], although one may also explore the alternative viewpoint, namely, through a modified gravity approach, in particular, using generalizations of the Einstein-Hilbert Lagrangian [4–7]. In fact, it is fundamental to understand how one may differentiate these modified theories of gravity from dark energy models. Note that generalizations of the action functional can be approached in several ways. For instance, prescriptions consist in replacing the linear scalar curvature term in the Einstein-Hilbert action by a function of the scalar curvature, $f(R)$ gravity [4], or by more general scalar invariants of the theory [5], or still considering a coupling between curvature and matter [6], a class of theories often termed higher-order gravity theories. Different approaches have been analysed in the literature: The metric formalism [4], which consists in varying the action with respect to $g^{\mu\nu}$. The Palatini formalism [8, 9], where the metric and the connections are treated as separate variables; and the metric-affine formalism, where the matter part of the action now depends and is varied with respect to the connection [9].

A novel approach to modified theories of gravity was recently proposed, consisting of adding to the metric Einstein-Hilbert Lagrangian an $f(\mathcal{R})$ term constructed à la Palatini [10]. It was shown that using the respective dynamically equivalent scalar-tensor representation, the theory can pass Solar System observational constraints even if the scalar field is very light. This implies the existence of a long-range scalar field, which is able to modify the cosmological [11] and galactic dynamics [12], but leaves the Solar System unaffected. Static and spherically symmetric solutions, in particular, wormhole solutions, were analysed in [13]. A generalization of the hybrid metric-Palatini gravitational theory was also presented [14], where the gravitational action is taken to depend on a general function of both the metric and Palatini curvature scalars. The dynamical equivalence with a non-minimally coupled bi-scalar field gravitational theory was proved and the evolution of cosmological solutions was studied using dynamical systems techniques.

---

*Electronic address: c.boehmer@ucl.ac.uk
†Electronic address: flobo@cii.fc.ul.pt
‡Electronic address: n.tamanini.11@ucl.ac.uk

In [15] the ghost analysis for these theories was recently performed showing that this biscalar generalization always presents either ghosts or tachyons. In fact, new types of second, fourth and sixth order derivative gravity theories were investigated and the hybrid metric-Palatini theory was singled out as a viable class of "hybrid" extensions of General Relativity.

The cosmological applications of this hybrid metric-Palatini gravitational theory were also further explored [11]. More specifically, cosmological solutions coming from its scalar-tensor representation of hybrid metric-Palatini gravity were found and criteria to obtain the cosmic acceleration were discussed. Furthermore, several classes of dynamical cosmological solutions, depending on the functional form of the effective scalar field potential, describing both accelerating and decelerating Universes were explicitly obtained. The cosmological perturbation equations were also derived and applied to uncover the nature of the propagating scalar degree of freedom and the signatures these models predict in the large-scale structure.

Relative to the galactic dynamics, the generalized virial theorem and certain astrophysical applications in the scalar-tensor representation of the hybrid metric-Palatini gravity were explored in [12]. More specifically, taking into account the relativistic collisionless Boltzmann equation, it was shown that the supplementary geometric terms in the gravitational field equations provide an effective contribution to the gravitational potential energy. It was also shown that the total virial mass is proportional to the effective mass associated with the new terms generated by the effective scalar field, and the baryonic mass. This fact demonstrates that the geometric origin in the generalized virial theorem may account for the well-known virial theorem mass discrepancy in clusters of galaxies. Thus, hybrid metric-Palatini gravity provides an effective alternative to the dark matter paradigm of present day cosmology and astrophysics.

In this work, in the cosmological context, motivated by the fact that the Einstein static Universe has always been of great interest in various gravitational theories, and inspired by the possibility that the universe may have started out in an asymptotically Einstein static state in the inflationary universe context [16], we analyse the Einstein static Universe and its stability in the hybrid metric-Palatini gravitational theory. In fact, the Einstein static universe has been analysed in a plethora of contexts, such as: in General Relativity with a non-constant pressure [17]; in brane world models [18]; in Einstein-Cartan theory [19]; in loop quantum cosmology [20]; in $f(R)$ modified theories of gravity [21–23]; in modified Gauss-Bonnet gravity [24]; in IR modified Hořava gravity [25]; and in massive gravity [26].

This present paper is outlined in the following manner: In Sec. II, we briefly review the formalism of the hybrid metric-Palatini gravitational theory, in particular the action and field equations, and in Sec. III present the respective modified Friedman equations. In Sec. IV, we consider linear homogeneous perturbations in the context of the Einstein static Universe in the above-mentioned hybrid metric-Palatini theory, and analyze the respective stability regions. In Sec. V, we summarize and discuss our results.

II. HYBRID METRIC-PALATINI GRAVITY: FORMALISM

The action for the hybrid metric-Palatini gravity is

$$S=\frac{1}{2\kappa^2}\int d^4x\sqrt{-g}\,[R+f(\mathcal{R})]+S_m, \tag{1}$$

where $\kappa^2\equiv 8\pi G$ and we set $c=1$. S_m is the matter action, R is the metric Einstein-Hilbert term, $\mathcal{R}\equiv g^{\mu\nu}\mathcal{R}_{\mu\nu}$ is the Palatini curvature. $\mathcal{R}_{\mu\nu}$ is defined in terms of an independent torsion-less connection $\hat\Gamma^\alpha_{\mu\nu}$ as

$$\mathcal{R}_{\mu\nu}\equiv\hat\Gamma^\alpha_{\mu\nu,\alpha}-\hat\Gamma^\alpha_{\mu\alpha,\nu}+\hat\Gamma^\alpha_{\alpha\lambda}\hat\Gamma^\lambda_{\mu\nu}-\hat\Gamma^\alpha_{\mu\lambda}\hat\Gamma^\lambda_{\alpha\nu}\,. \tag{2}$$

Varying the action (1) with respect to the metric, one obtains the following gravitational field equations

$$G_{\mu\nu}+F(\mathcal{R})\mathcal{R}_{\mu\nu}-\frac{1}{2}f(\mathcal{R})g_{\mu\nu}=\kappa^2T_{\mu\nu}\,, \tag{3}$$

where the matter stress-energy tensor is defined as

$$T_{\mu\nu}\equiv\frac{2}{\sqrt{-g}}\frac{\delta(\sqrt{-g}\mathcal{L}_m)}{\delta g^{\mu\nu}}\,. \tag{4}$$

The independent connection is compatible with the metric $F(\mathcal{R})g_{\mu\nu}$, which is conformal to $g_{\mu\nu}$, and where the conformal factor is given by $F(\mathcal{R})\equiv df(\mathcal{R})/d\mathcal{R}$. The latter considerations imply that

$$\begin{aligned}\mathcal{R}_{\mu\nu} \;=\; & R_{\mu\nu}+\frac{3}{2}\frac{1}{F^2(\mathcal{R})}F(\mathcal{R})_{,\mu}F(\mathcal{R})_{,\nu}\\ & -\frac{1}{F(\mathcal{R})}\nabla_\mu F(\mathcal{R})_{,\nu}-\frac{1}{2}\frac{1}{F(\mathcal{R})}g_{\mu\nu}\Box F(\mathcal{R})\,.\end{aligned} \tag{5}$$

Note that $\mathcal{R}$ can be obtained from the trace of the field equations (3), which yields

$$F(\mathcal{R})\mathcal{R}-2f(\mathcal{R})-R=\kappa^2T\,. \tag{6}$$

Introducing an auxiliary field, the hybrid metric-Palatini action (1) can be turned into a scalar-tensor theory given by the following action (we refer the reader to [10] for more details)

$$S=\frac{1}{2\kappa^2}\int d^4x\sqrt{-g}\,[R+\phi\mathcal{R}-V(\phi)]+S_m\,. \tag{7}$$

Varying this action with respect to the metric, the scalar ϕ and the connection yields the following field equations

$$R_{\mu\nu}+\phi\mathcal{R}_{\mu\nu}-\frac{1}{2}\left(R+\phi\mathcal{R}-V\right)g_{\mu\nu} \;=\; \kappa^2T_{\mu\nu}\,, \tag{8}$$

$$\mathcal{R}-V_\phi \;=\; 0\,, \tag{9}$$

$$\hat\nabla_\alpha\left(\sqrt{-g}\phi g^{\mu\nu}\right) \;=\; 0\,, \tag{10}$$

respectively. Note that the solution of Eq. (10) implies that the independent connection is the Levi-Civita connection of a metric $h_{\mu\nu} = \phi g_{\mu\nu}$. Thus we are dealing with a bi-metric theory and $\mathcal{R}_{\mu\nu}$ and $R_{\mu\nu}$ are related by

$$\mathcal{R}_{\mu\nu} = R_{\mu\nu} + \frac{3}{2\phi^2}\partial_\mu\phi\partial_\nu\phi - \frac{1}{\phi}\left(\nabla_\mu\nabla_\nu\phi + \frac{1}{2}g_{\mu\nu}\Box\phi\right) , \tag{11}$$

and consequently

$$\mathcal{R} = R + \frac{3}{2\phi^2}\partial_\mu\phi\partial^\mu\phi - \frac{3}{\phi}\Box\phi, \tag{12}$$

which can be used in the action (7) to eliminate the independent connection and obtain the following scalar-tensor representation [10]

$$\begin{aligned} S \;=\; & \frac{1}{2\kappa^2}\int d^4x\sqrt{-g}\left[(1+\phi)R + \frac{3}{2\phi}\partial_\mu\phi\partial^\mu\phi - V(\phi)\right] \\ & +S_m. \end{aligned} \tag{13}$$

It is important to note that this action differs fundamentally from the $w = -3/2$ Brans-Dicke theory due to the coupling of the scalar to the curvature. The trace of Eq. (8) yields $-R - \phi\mathcal{R} + 2V = \kappa^2 T$, and using Eq. (9), takes the following useful form

$$2V - \phi V_\phi = \kappa^2 T + R \ . \tag{14}$$

Now substituting Eq. (9) and Eq. (11) in Eq. (8), the metric field equation can be written as an effective Einstein field equation, i.e., $G_{\mu\nu} = \kappa^2 T^{\rm eff}_{\mu\nu}$, where the effective stress-energy tensor is given by

$$\begin{aligned} T^{\rm eff}_{\mu\nu} \;=\; & \frac{1}{1+\phi}\Big\{T_{\mu\nu} - \frac{1}{\kappa^2}\Big[\frac{1}{2}g_{\mu\nu}\left(V + 2\Box\phi\right) + \\ & \nabla_\mu\nabla_\nu\phi - \frac{3}{2\phi}\partial_\mu\phi\,\partial_\nu\phi + \frac{3}{4\phi}g_{\mu\nu}(\partial\phi)^2\Big]\Big\} . \end{aligned} \tag{15}$$

The scalar field is governed by the second-order evolution equation (we refer the reader to [10] for more details)

$$-\Box\phi + \frac{1}{2\phi}\partial_\mu\phi\partial^\mu\phi + \frac{\phi[2V - (1+\phi)V_\phi]}{3} = \frac{\phi\kappa^2}{3}T \, , \tag{16}$$

which is an effective Klein-Gordon equation. Equation (16) shows that the scalar field is dynamical, contrary to the Palatini case where $w = -3/2$. Thus, the theory is not affected by the microscopic instabilities that arise in Palatini models with infrared corrections.

III. EINSTEIN STATIC UNIVERSE AND MODIFIED FRIEDMANN EQUATIONS

A. Effective Friedmann equations

The cosmological applications of the hybrid metric-Palatini gravitational theory have been extensively explored [10, 11], in the scalar-tensor representation. In particular, it was shown that the accelerating expansion and, in general, any cosmological behaviour, strictly depends on the effective scalar field potential. However, it is rather important to stress that the scalar field has a purely geometric origin and describes further degrees of freedom of the gravitational field coming from extended theories of gravity.

Now, consider the Friedmann-Robertson-Walker (FRW) line element given by

$$ds^2 = -dt^2 + a^2(t)\left[\frac{dr^2}{1-Kr^2} + r^2(d^2\theta + \sin^2\theta d^2\phi)\right] \tag{17}$$

where $K = +1, 0, 1$ corresponds to a closed, flat, and open universe, respectively. For this case, the scalar curvature takes the following form

$$R = 6\left(2H^2 + \dot{H} + \frac{K}{a^2}\right) . \tag{18}$$

The effective Friedmann equations can be written in terms of the effective energy density and pressure, given by

$$3H^2 \;=\; \kappa^2\rho_{\rm eff} - \frac{3K}{a^2} \, , \tag{19}$$

$$\dot{H} \;=\; -\frac{\kappa^2}{2}\left(\rho_{\rm eff} + p_{\rm eff}\right) + \frac{K}{a^2} \, , \tag{20}$$

with the following relationships

$$\left(1+\phi\right)\kappa^2\rho_{\rm eff} \;=\; -\frac{3}{4\phi}\dot{\phi}^2 + \frac{1}{2}V(\phi) - 3H\dot{\phi} + \kappa^2\rho_m \, , \tag{21}$$

$$\left(1+\phi\right)\kappa^2 p_{\rm eff} \;=\; -\frac{3}{4\phi}\dot{\phi}^2 - \frac{1}{2}V(\phi) + \ddot{\phi} + 2H\dot{\phi} + \kappa^2 p_m \, . \tag{22}$$

The conservation equations for the matter component and the scalar field are

$$\dot{\rho}_m + 3H(\rho_m + p_m) \;=\; 0 \, , \tag{23}$$

$$\ddot{\phi} + 3H\dot{\phi} - \frac{\dot{\phi}^2}{2\phi} + \frac{1}{3}\phi R - \frac{1}{3}\phi V'(\phi) \;=\; 0 \, . \tag{24}$$

B. Einstein static Universe

For the Einstein static universe we choose $a = a_0 =$ const, so that $H = \dot{H} = 0$, and the curvature scalar reduces to $R = 6K/a_0^2$. Note that we do not specify K at this point. We also consider that the matter distribution obeys the following linear equation of state $p_m = w\rho_m$. In classical General Relativity, the presence of a positive cosmological constant allows us to solve the analogue equations of (19) and (20) under the staticity assumptions with $K = 1$. With the above considerations, the effective Friedmann equations (19) and (20) reduce to

$$\kappa^2\rho_{\rm eff} = \frac{3K}{a_0^2} \, , \tag{25}$$

$$\frac{\kappa^2}{2}\left(\rho_{\rm eff} + p_{\rm eff}\right) = \frac{K}{a_0^2} \, , \tag{26}$$

respectively, which imposes the following condition on the distribution of matter

$$\rho_{\text{eff}} + 3p_{\text{eff}} = 0\,. \tag{27}$$

As a first approach, let us now assume that $\phi = \phi_0 =$ const, then the condition on the effective matter equation implies

$$\kappa^2 \rho_m (1 + 3w) = V(\phi_0)\,, \tag{28}$$

and the modified Klein-Gordon equation (24) reduces to the following expression

$$\frac{6K}{a_0^2} = V'(\phi_0)\,. \tag{29}$$

Therefore, we can express ρ_m and a_0 in terms of ϕ_0 and the potential V. Substituting back into one of the two field equations, we can find an implicit equation which fixes the value of the field ϕ_0 by

$$1 + \phi_0 = \frac{V(\phi_0)}{V'(\phi_0)} \frac{3(1+w)}{(1+3w)}\,. \tag{30}$$

Depending on the functional form of the potential, it may be possible to solve for ϕ_0 explicitly.

One should also note at this point that $K = -1$ is a possible parameter for an Einstein static universe, in the context of the hybrid metric-Palatini gravitational theory. Provided $V'(\phi_0) < 0$, one can find the scale factor a_0 using Eq. (29).

IV. STABILITY ANALYSIS OF THE EINSTEIN STATIC UNIVERSE

It is interesting that the Einstein static universe has been revived as the asymptotic origin of an emergent universe, more specifically as an inflationary cosmology without a singularity [16]. Despite the fact the positive curvature is negligible at late times, its role is crucial during the the early universe. In the latter context, it allows these cosmologies to inflate and later reheat to a hot big-bang epoch. In fact, these cosmological models possess attractive features such as the absence of a singularity, of an 'initial time', of the horizon problem, and the quantum regime can even be avoided.

In order to study the stability of the Einstein universe, we now consider the following homogeneous perturbations

$$a = a_0 + \delta a(t)\,, \tag{31}$$

$$\phi = \phi_0 + \delta\phi(t)\,, \tag{32}$$

$$\rho_m = \rho_m^{(0)} + \delta\rho_m(t)\,, \tag{33}$$

where we only consider adiabatic perturbations $\delta p_m(t) = w\,\delta\rho_m(t)$. When perturbing the field equations (19) and (20), and the evolution equations for the matter (23) and (24), we will only consider linear perturbations and neglect all terms of higher order.

Let us begin with equation (23). Perturbing to linear order provides

$$\dot{\delta\rho_m} + 3(1+w)\frac{\rho_m^{(0)}}{a_0}\dot{\delta a} = 0\,, \tag{34}$$

which can be integrated to yield the following relationship

$$\delta\rho_m = -3(1+w)\frac{\rho_m^{(0)}}{a_0}\delta a\,, \tag{35}$$

which can be used to substitute the matter perturbation.

Note that we have three independent equations for the three quantities δa, $\delta\phi$ and $\delta\rho_m$. Thus, we have two more equations to solve, these are equations (20) and (24). We are now linearising these with respect to the perturbations and substitute (35) and the background solutions (25)–(30) into the field equations (20) and (24). This results in two second order ordinary differential equations. It is useful to write the result in the following form

$$\begin{pmatrix} \ddot{\delta a} \\ \ddot{\delta\phi} \end{pmatrix} = \begin{pmatrix} M_{11} & M_{12} \\ M_{21} & M_{22} \end{pmatrix} \begin{pmatrix} \delta a \\ \delta\phi \end{pmatrix}\,, \tag{36}$$

where the components of the matrix $\mathbf{M}$ are given by

$$M_{11} = \frac{3V(\phi_0)(1+w)(-1+3w) + 2V'(\phi_0)(1+3w)}{6(1+3w)}\,, \tag{37}$$

$$M_{12} = \sqrt{K}\frac{-3V(\phi_0)V''(\phi_0)(1+w) + V'(\phi_0)[V'(\phi_0) + V''(\phi_0)](1+3w)}{\sqrt{6}V'(\phi_0)^{3/2}(1+3w)}\,, \tag{38}$$

$$M_{21} = \frac{V(\phi_0)V'(\phi_0)^{3/2}(1+w)(-1+3w)[V'(\phi_0)(1+3w) - 3V(\phi_0)(1+w)]}{\sqrt{K}\sqrt{6}V'(\phi_0)^2(1+3w)^2}\,, \tag{39}$$

$$M_{22} = \frac{[V'(\phi_0)(1+3w) - 3V(\phi_0)(1+w)][-3V(\phi_0)V''(\phi_0)(1+w) + V'(\phi_0)^2(1+3w)]}{3V'(\phi_0)^2(1+3w)^2}\,. \tag{40}$$

It is more convenient to work with a first order than a second order system of equations. Thus, introducing two new variables, $\delta b = \dot{\delta a}$ and $\delta\psi = \dot{\delta\phi}$, Eq. (36) can be rewritten as

$$\begin{pmatrix} \dot{\delta a} \\ \dot{\delta\phi} \\ \dot{\delta b} \\ \dot{\delta\psi} \end{pmatrix} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ M_{11} & M_{12} & 0 & 0 \\ M_{21} & M_{22} & 0 & 0 \end{pmatrix} \begin{pmatrix} \delta b \\ \delta\psi \\ \delta a \\ \delta\phi \end{pmatrix} , \tag{41}$$

This last equation is of the form $\dot{\boldsymbol{v}} = \mathbf{A}\boldsymbol{v}$, and the eigenvalues of the 4×4 matrix $\mathbf{A}$ determine the stability or instability properties of the perturbations. Let us denote the two eigenvalues of $\mathbf{M}$ as λ_1 and λ_2, then the four eigenvalues of $\mathbf{A}$ are simply given by $\pm\sqrt{\lambda_1}$ and $\pm\sqrt{\lambda_2}$. We find

$$\lambda_1 = \frac{1}{2}\Big(M_{11} + M_{22} + \sqrt{(M_{11} - M_{22})^2 + 4M_{12}M_{21}}\Big) , \tag{42}$$

$$\lambda_2 = \frac{1}{2}\Big(M_{11} + M_{22} - \sqrt{(M_{11} - M_{22})^2 + 4M_{12}M_{21}}\Big) . \tag{43}$$

In order for stable perturbation to exist, we need to require that both eigenvalues are purely real, and additionally $\lambda_1 < 0$ and $\lambda_2 < 0$. The reality condition entails

$$(M_{11} - M_{22})^2 + 4M_{12}M_{21} > 0 , \tag{44}$$

while the negativity condition implies

$$M_{11} + M_{22} < -\sqrt{(M_{11} - M_{22})^2 + 4M_{12}M_{21}} , \tag{45}$$

$$(M_{11} + M_{22})^2 < (M_{11} - M_{22})^2 + 4M_{12}M_{21} . \tag{46}$$

The latter conditions can be rewritten as

$$(M_{11} + M_{22})^2 - (M_{11} - M_{22})^2 < 4M_{12}M_{21} , \tag{47}$$

$$4M_{11}M_{22} < 4M_{12}M_{21} , \tag{48}$$

$$M_{11}M_{22} - M_{12}M_{21} < 0 , \tag{49}$$

$$\det \mathbf{M} < 0 . \tag{50}$$

Therefore we have the necessary condition $\det \mathbf{M} < 0$.

Taking both conditions together, namely the reality and the negativity conditions, we require

$$(M_{11} - M_{22})^2 + 4M_{12}M_{21} > 0 \tag{51}$$

$$M_{11} + M_{22} < -\sqrt{(M_{11} - M_{22})^2 + 4M_{12}M_{21}} . \tag{52}$$

As our expression for the components of $\mathbf{M}$ are quite involved, it is expected that those inequalities are very complicated when expressed in terms of $V(\phi_0), V'(\phi_0), V''(\phi_0)$ and w. However, we can indicate graphically the parameter space where the conditions are satisfied. Firstly, we note that we can always rescale the values of the potential at ϕ_0, therefore we will set $V(\phi_0) = 1$. For a fixed value of w, the two conditions now depend on the two values $V'(\phi_0)$ and $V''(\phi_0)$. We plot the relevant regions in the Figures 1–3.

In Figures 1–3 the shaded regions represent the parameter space where homogeneous perturbations of the Einstein static universe are stable. In each figure the equation of state parameter w is fixed, we chose the six representative values $w = \{1/2, 1/5, 0, -1/5, -3/5, -4/5\}$, respectively. Note that a wide variety of stability regions exist for a range of positive and negative values of $V'(\phi_0)$. One could have attempted a three dimensional plot with axes $V'(\phi_0), V''(\phi_0), w$, however, this gives a very complicated figure of little value. We should emphasise that these stability figures have to be seen in context with the background solutions (25)–(30). For instance, we require a positive effective energy density in (25) and a positive value for a_0 in (29), both of which are independent of the stability of the perturbations. We should also note that once a specific potential has been chosen, the terms $V'(\phi_0)$ and $V''(\phi_0)$ are not necessarily independent anymore and care has to be taken when interpreting the figures.

V. SUMMARY AND DISCUSSION

In this work, we analysed the stability of the Einstein static Universe by considering linear homogeneous perturbations in the respective dynamically equivalent scalar-tensor representation of hybrid metric-Palatini gravity. Considering a linear equation of state parameter for the matter distribution, the stability regions of the Einstein static universe were parametrized by the first and second derivatives of the scalar potential, and it was shown that a large class of stable solutions exists in the respective parameter space. The stability regions were considered by requiring positive values for the effective energy density and for the scale factor a_0, both of which are independent of the stability of the perturbations.

It is interesting to discuss a few specific choices for the potential $V(\phi)$ and search for stable static solutions. For instance, consider the exponential potential given by

$$V(\phi) = V_0\, e^{-\lambda\phi} , \tag{53}$$

where V_0 and λ are two positive constants. In an open universe ($K = -1$) a static solution is given by

$$a_0 = \sqrt{\frac{6}{A\,\lambda}}\, \exp\left[-\frac{\lambda + 3(\lambda+1)w + 3}{6w + 2}\right] , \tag{54}$$

$$\rho_0 = \frac{A}{\kappa^2(3w+1)}\, \exp\left[\lambda + \frac{2}{3w+1} + 1\right] , \tag{55}$$

$$\phi_0 = -\frac{\lambda + 3(\lambda+1)w + 3}{\lambda(1+3w)} , \tag{56}$$

where $w > -1/3$ in order to have $\rho_0 > 0$. This is quite interesting as the effects of hybrid metric-Palatini gravity are sufficiently strong to allow for a static solution in the open case. Unfortunately this solution is always unstable for $w > -1/3$. This is in line with the point made above,



FIG. 1: Left: Stability regions for $w = 1/2$. Right: Stability regions for $w = 1/5$. The shaded regions of the figures represent the parameter space where homogeneous perturbations of the Einstein static universe are stable. In both cases one cannot find stable solutions unless $V'(\phi_0) < 0$.



FIG. 2: Left: Stability regions for $w = 0$. Right: Stability regions for $w = -1/5$. As in the previous figure, the shaded regions of the figures represent the parameter space where homogeneous perturbations of the Einstein static universe are stable, and in both cases the stability regions exist for $V'(\phi_0) < 0$.

namely that the stability regions have to be interpreted using specific background solutions.

Let us now set $K = 1$. Recall that in General Relativity the Einstein static universe only exists if $K = 1$. However, for this case, there are no real and positive solutions for a_0 in this closed universe with the exponential potential (53). This follows from (29), the exponential potential has negative derivative and for positive K the scale factor a_0 cannot be real. This indicates that one should consider a different potential.

As a second specific potential, consider the quadratic potential given by

$$V(\phi) = \frac{1}{2}m^2\phi^2\,, \tag{57}$$

where m is a positive constant scalar field mass. A static



FIG. 3: Left: Stability regions for $w = -3/5$. Right: Stability regions for $w = -4/5$. The shaded regions of the figures represent the parameter space where homogeneous perturbations of the Einstein static universe are stable. The equation of state parameter now satisfies $w < -1/3$ and one can find stable solutions for a wide variety of positive values of $V'(\phi_0)$.

solution for this potential is given by

$$a_0 = \sqrt{\frac{3\,k\,(1-3w)}{A\,(6w+2)}}\,, \tag{58}$$

$$\rho_0 = \frac{4\,A\,(3w+1)}{\kappa^2\,(1-3w)^2}\,, \tag{59}$$

$$\phi_0 = \frac{6w+2}{1-3w}\,. \tag{60}$$

As in the previous case, the background solution will place constraints on the permissible parameter range. In an open universe ($K = -1$) we must require $w > 1/3$ in order to have a_0 and ρ_0 both real and positive, while in a closed universe ($K = 1$) we need $-1/3 < w < 1/3$. The potential satisfies $V''(\phi) = m^2$ and is thus strictly positive. Combining this with the condition on w, we conclude that the open universe solution is always unstable. On the other hand, the closed universe is stable for every allowed value of w.

Despite concentrating the analysis in this work to the stability of the Einstein static Universe by considering linear homogeneous perturbations, which has yielded interesting results and has presented a richer stability/instability structure that in General Relativity, it is of interest to extend our results to inhomogeneous perturbations. Work along these lines is presently underway.

Acknowledgments

FSNL acknowledges financial support of the Fundação para a Ciência e Tecnologia through the grants CERN/FP/123615/2011 and CERN/FP/123618/2011.

---

[1] S. Perlmutter *et al.* [Supernova Cosmology Project Collaboration], Astrophys. J. **517**, 565 (1999); A. G. Riess *et al.* [Supernova Search Team Collaboration], Astron. J. **116**, 1009 (1998). A. G. Riess *et al.* [Supernova Search Team Collaboration], Astrophys. J. **607**, 665 (2004).

[2] P. A. R. Ade et al., Planck 2013 results. I, arXiv: 1303.5062[astro-ph) (2013); P. A. R. Ade et al., Planck 2013 results. XVI, arXiv: 1303.5076[astro-ph] (2013); P.A.R. Ade et al., Planck 2013 results. XXII, arXiv: 1303.5082[astro-ph] (2013); P. A. R. Ade et al., Planck 2013 Results. XXIV, arXiv:1303.5084[astro-ph] (2013); P. A. R. Ade et al., Planck 2013 Results. XXVI, arXiv:1303.5086 (astro-ph) (2013).

[3] E. J. Copeland, M. Sami and S. Tsujikawa, Int. J. Mod. Phys. D **15**, 1753 (2006).

[4] S. Capozziello, Int. J. Mod. Phys. **D 11**, 483 (2002); S. M. Carroll, V. Duvvuri, M. Trodden and M. S. Turner, Phys. Rev. D **70**, 043528 (2004); S. Nojiri and S.D. Odintsov, Int.J.Geom.Meth.Mod.Phys. **4**, 115 (2007); S. Capozziello and M. Francaviglia, Gen. Rel. Grav. **40**, 357 (2008); T. P. Sotiriou and V. Faraoni, Rev. Mod. Phys. **82**, 451 (2010); A. De Felice and S. Tsujikawa, Living Rev. Rel. **13**, 3 (2010); F. S. N. Lobo, arXiv:0807.1640 [gr-qc]; S. Capozziello and M. De Laurentis, Phys. Rept. **509**, 167 (2011); S. 'i. Nojiri and S. D. Odintsov, Phys. Rept. **505**, 59 (2011).

[5] S. Nojiri, S. D. Odintsov and M. Sasaki, Phys. Rev. D **71**, 123509 (2005).

[6] O. Bertolami, C. G. Boehmer, T. Harko and F. S. N. Lobo, Phys. Rev. D **75**, 104016 (2007); O. Bertolami, F. S. N. Lobo and J. Paramos, Phys. Rev. D **78**, 064036 (2008); O. Bertolami, J. Paramos, T. Harko and F. S. N. Lobo, arXiv:0811.2876 [gr-qc]; T. Harko, F. S. N. Lobo, S. 'i. Nojiri and S. D. Odintsov, Phys. Rev. D **84**, 024020 (2011); T. Harko and F. S. N. Lobo, Eur. Phys. J. C **70**, 373 (2010); T. Harko, F. S. N. Lobo and O. Minazzoli, Phys. Rev. D **87**, 047501 (2013); Z. Haghani, T. Harko, F. S. N. Lobo, H. R. Sepangi and S. Shahidi, arXiv:1304.5957 [gr-qc]; S. D. Odintsov and D. Sáez-Gómez, arXiv:1304.5411 [gr-qc].

[7] C. G. Boehmer and N. Tamanini, arXiv:1301.5471 [gr-qc].

[8] M. Ferraris, M. Francaviglia and I. Volovich, gr-qc/9303007 (1993); D. N. Vollick, Phys. Rev. **D68**, 063510 (2003); E. E. Flanagan, Class. Quant. Grav. **21**, 417 (2003); X. H. Meng and P. Wang, Phys. Lett. **B584**, 1 (2004); B. Li and M. C. Chu, Phys. Rev. **D74**, 104010 (2006); N. J. Poplawski, Phys. Rev. **D74**, 084032 (2006); B. Li, K. C. Chan and M. C. Chu, Phys. Rev. **D76**, 024002 (2007); B. Li, J. D. Barrow and D. F. Mota, Phys. Rev. D **76**, 104047 (2007); A. Iglesias, N. Kaloper, A. Padilla and M. Park, Phys. Rev. D **76**, 104001 (2007); G. J. Olmo, Int. J. Mod. Phys. D **20**, 413-462 (2011).

[9] T. P. Sotiriou and S. Liberati, Annals Phys. **322**, 935 (2007).

[10] T. Harko, T. S. Koivisto, F. S. N. Lobo and G. J. Olmo, Phys. Rev. D **85**, 084016 (2012).

[11] S. Capozziello, T. Harko, T. S. Koivisto, F. S. N. Lobo and G. J. Olmo, JCAP **1304**, 011 (2013) [arXiv:1209.2895 [gr-qc]].

[12] S. Capozziello, T. Harko, T. S. Koivisto, F. S. N. Lobo and G. J. Olmo, arXiv:1212.5817 [physics.gen-ph].

[13] S. Capozziello, T. Harko, T. S. Koivisto, F. S. N. Lobo and G. J. Olmo, Phys. Rev. D **86**, 127504 (2012) [arXiv:1209.5862 [gr-qc]].

[14] N. Tamanini and C. G. Boehmer, Phys. Rev. D **87** (2013) 084031; E. E. Flanagan, Class. Quant. Grav. **21** (2003) 417.

[15] T. S. Koivisto and N. Tamanini, arXiv:1304.3607 [gr-qc].

[16] G. F. R. Ellis and R. Maartens, Class. Quant. Grav. **21** (2004) 223.

[17] A. Ibrahim and Y. Nutku, Gen. Rel. Grav. **7** (1976) 949; C. G. Böhmer, [arXiv:gr-qc/0308057]; C. G. Böhmer, Gen. Rel. Grav. **36** (2004) 1039.

[18] L. A. Gergely and R. Maartens, Class. Quant. Grav. **19** (2002) 213; A. Gruppuso, E. Roessl and M. Shaposhnikov, JHEP **0408**, 011 (2004); S. S. Seahra, C. Clarkson and R. Maartens, Class. Quant. Grav. **22** (2005) L91; C. Clarkson and S. S. Seahra, Class. Quant. Grav. **22** (2005) 3653.

[19] C. G. Böhmer, Class. Quant. Grav. **21** (2004) 1119.

[20] D. J. Mulryne, R. Tavakol, J. E. Lidsey and G. F. R. Ellis, Phys. Rev. **D71** (2005) 123512; L. Parisi, M. Bruni, R. Maartens and K. Vandersloot, Class. Quant. Grav. **24** (2007) 6243.

[21] C. G. Böhmer, L. Hollenstein and F. S. N. Lobo, Phys. Rev. **D76** (2007) 084005.

[22] R. Goswami, N. Goheer and P. K. S. Dunsby, Phys. Rev. **D78** (2008) 044011; N. Goheer, R. Goswami and P. K. S. Dunsby, Class. Quant. Grav. **26** (2009) 105003 arXiv:0809.5247 [gr-qc].

[23] S. S. Seahra and C. G. Böhmer, Phys. Rev. D **79** (2009) 064009 arXiv:0901.0892 [gr-qc].

[24] C. G. Böhmer and F. S. N. Lobo, Phys. Rev. **D79** (2009) 067504.

[25] C. G. Boehmer and F. S. N. Lobo, Eur. Phys. J. C **70**, 1111 (2010); P. Wu and H. W. Yu, Phys. Rev. D **81** (2010) 103522; R. Canonico and L. Parisi, Phys. Rev. D **82** (2010) 064005.

[26] L. Parisi, N. Radicella and G. Vilasi, Phys. Rev. D **86** (2012) 024035.